



Review

The Evi5 family in cellular physiology and pathology

Yi Shan Lim^a, Bor Luen Tang^{a,b,*}^a Department of Biochemistry, Yong Loo Lin School of Medicine, National University Health System, Singapore^b NUS Graduate School of Integrative Sciences and Engineering, National University of Singapore, 8 Medical Drive, Singapore 117597, Singapore

ARTICLE INFO

Article history:

Received 25 February 2013

Revised 23 April 2013

Accepted 28 April 2013

Available online 10 May 2013

Edited by Lukas Huber

Keywords:

Cell cycle

Evi5

GTPase Activating Protein (GAP)

Rab

ABSTRACT

The Ecotropic viral integration site 5 (Evi5) and Evi5-like (Evi5L) belong to a small subfamily of the Tre-2/Bub2/Cdc16 (TBC) domain-containing proteins with enigmatically divergent roles as modulators of cell cycle progression, cytokinesis, and cellular membrane traffic. First recognized as a potential oncogene and a cell cycle regulator, Evi5 acts as a GTPase Activating Protein (GAP) for Rab11 in cytokinesis. On the other hand, its homologue Evi5L has Rab-GAP activity towards Rab10 as well as Rab23, and has been implicated in primary cilia formation. Recent genetic susceptibility analysis points to Evi5 as an important factor in susceptibility to multiple sclerosis. We discuss below the myriad of cellular functions exhibited by the Evi5 family members, and their associations with disease conditions.

© 2013 Federation of European Biochemical Societies. Published by Elsevier B.V. All rights reserved.

1. Introduction

The mouse gene locus Ecotropic viral integration site 5 (Evi5) was first identified as a common site for retroviral integration at mouse chromosome 5 in T-cell lymphomas of recombinant inbred AKXD mice with high incidences of lymphoma [1]. Pro-viral integrations at the Evi5 locus disrupt a gene [2] – located 18 kb upstream of another common viral integration site, Growth factor independence 1 (*Gfi1*) (which itself encodes a zinc finger-containing transcription factor involved in interleukin-2 signaling). The human orthologue was separately identified as part of the constitutional, reciprocal, t(1;10)(p22q21) translocation breakpoint in a patient with stage 4S neuroblastoma. The corresponding gene on chromosome 1p22, also named NB4S [3], encodes a 7.5 kbp transcript with an 810 amino acid open reading frame expressed in a wide variety of tissues. A paralogue, known as Evi5 homologue [4] or Evi5-like [5], shares about 70% identity to Evi5 and contains the same domain structure, is located on mouse chromosome 8 and human chromosome 19 respectively.

Human and mouse Evi5 and Evi5-like encodes multiple alternatively spliced transcripts. The longest polypeptide encoded by these two proteins share two prominent domain features—a Tre-2/Bub2/Cdc16 (TBC) domain [6] at the N-terminal half, as well as a coiled-coil region that bears homology to Structural Maintenance

of Chromosomes (SMC) family of ATPases [7] at the C-terminal half (Fig. 1). TBC domains are commonly associated with a functional class of proteins that act as GTPase Activating Proteins (GAPs) for the Rab family GTPases that regulate membrane traffic [8]. SMC proteins, on the other hand, are an evolutionarily conserved group of molecules that are important in maintaining genome integrity [7,9]. In eukaryotes, they are core components of the condensin and cohesin complexes responsible for regulating chromosomal condensation, pairing and segregation [10]. The physiological functions of Evi5 are only beginning to be unraveled, and characterization of Evi5's localization and activities indicate that it may serve different functions during the course of cell division by engaging different interacting partners. Furthermore, Evi5L may be functionally distinct from Evi5 in spite of their high degree of homology. We discuss in the paragraphs below, previously known and recently revealed cellular activities exhibited by the Evi5 family members, as well as their implicated or predicted involvements in disease pathology.

2. Evi5 as a modulator of events during cell division

The initial reports describing the discovery of Evi5 shed little light on its possible function other than hinting that the ubiquitously expressed gene product, or truncated forms of it, could contribute to oncogenesis, which is unsurprising given the link between TBC domains and cell cycle regulation [2,3]. A subsequent report indicated that Evi5 has a putative nuclear localizing signal and exhibited a diffused nucleoplasm staining [11]. Interestingly, Evi5 is a phosphoprotein and has a prominent centrosomal

* Corresponding author at: NUS Graduate School of Integrative Sciences and Engineering, National University of Singapore, 8 Medical Drive, Singapore 117597, Singapore. Fax: +65 6779 1453.

E-mail address: bchtlb@nus.edu.sg (B.L. Tang).




Fig. 1. Schematic diagram of human TBC1D20, Evi5 and Evi5L. Protein sequence analyses of complete sequences of human TBC1D20 (an example of a known RabGAP that accelerates hydrolysis of Rab1), human Evi5 and human Evi5L, were carried out using two public domain software, Interpro (<http://www.ebi.ac.uk/interpro/>) protein sequence analysis and classification, and PSORT II prediction (<http://psort.hgc.jp/form2.html>). Boxes and arrows mark out the relative positions of TBC domains and the coiled-coil domains. The numbers along the protein sequence depict the relative amino acid positions of where the respective regions begin. The multiple alignment analysis was performed by ClustalW (<http://embnet.vital-it.ch/software/ClustalW.html>). Green highlighting marks residues that are identical between the TBC domains of TBC1D20, Evi5 and Evi5L, pink highlights refer to conserved substitutions, while yellow highlights the semi-conserved substitutions.

localization in interphase cells. It also co-precipitates with both α - and γ -tubulin. This differential subcellular localization patterns indicate that Evi5 could potentially act in different compartments of the cell.

Further work revealed that Evi5 undergoes several post-translational modifications, and has a dynamic association with mitotic structures during the M phase [12]. It decorates the mitotic spindle through anaphase and remains within the mid-zone and mid-body until the completion of cytokinesis. The protein is phosphorylated in early mitosis and undergoes proteolytic cleavage during late mitosis and cytokinesis. Silencing of Evi5 resulted in a multinucleate phenotype, suggesting a role for Evi5 in the completion of cytokinesis. In this regard, it is intriguing that Evi5 was shown to associate with the Chromosomal Passenger Complex (CPC) consisting of Aurora B kinase, Inner Centromere Proteins (INCENP), survivin and borealin [12]. The CPC acts in multiple processes during mitosis that safeguards genomic integrity during cell division. During cytokinesis, while daughter cells are still connected by chromosome bridges, the CPC limits chromosomal damage by imposing an abscission delay [13]. A recent report suggests that the CPC controls abscission timing through Aurora B kinase mediated-phosphorylation and inhibition of the vacuolar-sorting protein Snf7 component of the Endosomal Sorting Complex Required for Transport-III (ESCRT-III) polymerization and membrane association [14]. The exact role of Evi5 in cytokinesis, and whether it acts in this regard via its association with the CPC, is still unknown. On the other hand, another hypothesis proposed is that Evi5 exerts its effect in cytokinesis via its control of Rab11 activity [15,16]. This shall be discussed in more detail below.

Evi5 has also another implicated role in the earlier phases of the cell cycle – as a regulator of cyclin accumulation via its effects on

the stability of Early mitotic inhibitor 1 (Emi1) accumulation [4]. Emi1 is an inhibitor of anaphase promoting complex/cyclosome (APC/C) complex, and together with either of its activators Cdc20 or Cdh1, sustain levels of APC/C's cyclin substrates [17,18]. Even in the absence of transcription factor E2F activation and basal levels of cyclin A transcription, Emi1's overexpression is apparently sufficient to drive S phase by stabilizing cyclin A [18]. Before mitosis, at the G1/S phase and throughout the S phase, Evi5 binds to a site adjacent to Emi1's degron, blocking degron phosphorylation by Polo-like kinases (Plks) [4], particularly Plk1 [19], thereby allowing the accumulation of Cyclin A for the progression of S phase [20]. Upon the onset of mitosis, phosphorylation of Emi1 leads to degradation by the Skp, Cullin, F-box (Fb) containing protein (SCF) E3 ubiquitin ligase complex, one of which consists of a β -transducin repeat containing protein (β TrCP)/Fbw1a [21]. Finally, phosphorylated Evi5 and Emi1 are then destined for ubiquitination and degradation by the 26S proteasome. Hence, Evi5 effectively maintains Emi1 levels in the S/G2 phase. Evi5 silencing induces aberrant and untimely degradation of Emi1, resulting in a sequence of events ranging from premature APC/C activation and cyclin destruction, culminating in cell-cycle arrest and mitotic catastrophe. Taken together, by acting through different interacting partners, Evi5 is able to exert varying influences on cell cycle progression (Fig. 2).

3. Evi5 as a Rab11 GAP

The Rab family of small GTPases regulate a myriad of membrane transport processes in the eukaryotic cell [22,23] via their engagement of effector molecules such as motors, tethering factors, coat proteins and membrane fusion factors [24–27] in their active GTP-bound form. Two classes of proteins principally




Fig. 2. Evi5's role in early stages of the cell cycle. Evi5 is a regulator of cyclin accumulation via its effects on the stability of the Early mitotic inhibitor 1 (Emi1) [4], the latter being an inhibitor of anaphase promoting complex/cyclosome (APC/C) complex. Before mitosis, at the G1/S phase and throughout the S phase, Evi5 binds to a site adjacent to Emi1's degron, blocking its phosphorylation by Polo-like kinase 1 (Plk1), thereby allowing the accumulation of Cyclin A for the progression into S phase. Cyclin B/Cdk1 appears to "prime" or enhance the phosphorylation of Emi1 by Plk1 [19]. Plk1 is also able to stimulate the phosphorylation of Evi5. Upon the onset of mitosis, Emi1 levels are regulated via degradation by the Skp, Cullin, F-box (Fb) containing protein (SCF) and β -transducin repeat containing protein (β TrCP) E3 ubiquitin ligase complex [21]. Phosphorylated Evi5 and Emi1 are then destined for ubiquitination and degradation by the 26S proteasome. See text for relevant references and sources.

regulate the GTP-binding status of Rabs, namely the Guanine nucleotide Exchange Factors (GEFs) and GTPase Activating Proteins [28]. GEFs catalyze the GTP–GDP exchange of the Ras superfamily of small GTPases to effectively activate them [29]. A small number of Rab GEFs since identified have been found to be largely structurally unrelated to each other. However, certain Rab GEFs do form small functional subfamilies that are distinguishable by signature domains such as Vacuolar protein sorting 9 (Vps9) [30] and differentially expressed in normal and neoplastic cells (DENN) [31] domains.

The intrinsic GTP hydrolysis activity of Rab proteins is usually very low, and requires enzymatic activity induced by their respective Rab GAP proteins. GAPs are important in enhancing hydrolysis of the GTP-bound Rab, thereby switching activated Rabs to the inactive state [32]. These proteins therefore serve to restrict spatial and temporal overlap of Rabs, by mediating sequential steps of membrane transport [33]. Compared to GEFs, known Rab GAPs are somewhat less diverse, with Tre-2/Bub2/Cdc16 (TBC) domain-containing proteins expected to function as a Rab GAP [8,28]. The TBC domain was first recognized as a domain that was conserved among the *tre-2* oncogene product and yeast cell cycle regulators Bub2 and Cdc16, beginning with the cloning of mammalian TBC1 [34], has to date been discovered in more than 40 other proteins in humans and mice [8]. The GAP of Ypt6p (Gyp6p), the first Rab GAP, was cloned from yeast [35]. Several others, including those from mammals acting principally on Rab5 and Rab6, were subsequently identified [36–40]. A yeast 2-hybrid-based screen and subsequent in vitro Rab GAP activity assays identified seven TBC domain-containing proteins that interact with specific Rabs, of which a fraction of these proteins exhibit significant Rab GAP activity [5]. Although the TBC domain is proposed to be important for Rab interaction and best known as the Rab-GAP domain, a recent protein interaction screen for Rab effectors (i.e. which binds to the activated form of Rabs) identified three known GAPs, namely, mKIAA1055/TBC1D2B (Rab22-binding protein), GAPCenA/TBC1D11 (Rab36-binding protein) and centaurin β 2 (Rab35-binding protein, which does not contain a TBC domain). Notably, all these three GAP proteins appear to interact with their specific Rabs via a domain other than the TBC domain [41].

Incidentally, two reports that were published in 2007 both demonstrated Evi5's interaction with members of the Rab11 family, also drew differing conclusions with regards to Evi5's GAP activity towards Rab11. The earlier paper by Westlake and

colleagues showed through genetic and biochemical interaction assays that Evi5 binds Rab11a and Rab11b in a GTP-dependent manner, but exhibited no GAP activity towards Rab11 in in vitro assays [42]. Despite that, endogenous Evi5 co-localizes with Rab11 in a complex in vivo and overexpression of Rab11 redistributes centrosomal cytoplasmic-localized Evi5 to Rab11-positive recycling endosomes, in U2OS cells, thus suggesting a relationship between Rab11 and Evi5. Also, Rab11 effector proteins such as Rab11 family interacting protein 3 (Rab11-FIP3), which bind to active, GTP-bound Rab11, could compete with Evi5 for Rab11 binding. Another paper by Dabbeek et al. also found that during interphase in the HeLa cell model, Rab11 and Evi5 co-localized at the pericentriolar region. Furthermore, Evi5 was shown to interact with Rab11 in a GTP-enhanced manner via its TBC domain. Contrary to the previous report, these authors confirmed that Evi5 exhibited specific Rab GAP activity in vitro towards Rab11 (in comparison with similarly endosome-localized Rab4) [43].

4. Evi5's association with the Rab11 subfamily of Rab GTPases

The Rab11 subfamily (Rab11a, Rab11b and Rab25) and their effectors [44] mediate trafficking steps at and from the recycling endosome [45,46] of a large variety of molecules such as ligand transporters [47,48], ion channels [49–51] signal receptors [52–58], cell adhesion molecules [59,60] and others [61]. Endosomal recycling activities mediated by the Rab11 subfamily are key events in various physiological processes such as regulation of cell polarity [62], cell fate determination [63], tissue morphogenesis [64,65], primary ciliogenesis [66], autophagy [67], membrane trafficking and cytokinesis [15,16,68,69], and cell migration [70,71]. Disruption of Rab11-mediated traffic could underlie pathologies associated with neurodegenerative disorders such as Huntington's disease [72,73], while aberrant Rab11 family-mediated traffic have been associated with cancer cell migration and invasion [60,74].

Evi5's interaction with Rab11 suggests that it could potentially have a wide range of physiological and developmental roles, many of which still lack elucidation. Foremost, Evi5 may play an important role in the regulation of abscission via Rab11 and Rab11-FIP3, because silencing of Evi5 by RNA interference disrupted localization of Rab11-FIP3 complexes at abscission sites of all randomly selected cells undergoing final stages of cytokinesis [42]. During cell division, Rab11-FIP3 has been implicated in the membrane delivery from endosomal recycling compartments to sites of membrane

insertion (Fig. 3). Detailed analysis of the roles played by Evi5 during cytokinesis could potentially be complicated by its apparent involvement in the earlier stages of the cell cycle, as discussed in the previous section. The understanding Evi5 and Rab11's possible role in cell migration has recently been further deepened. An RNAi-based screen for Rab GAPs involved in collective cell migration of *Drosophila melanogaster* border cells (BCs), which has been known to be dependent on protein trafficking, identified the fly orthologue of Evi5 as a key regulator of the BC migration [75]. Evi5 acts through Rab11, functioning as a Rab GAP for Rab11. In BC migration towards the oocyte, polarized spatial localization of receptor tyrosine kinase (RTK) activity at the leading edge is required. Moreover, an earlier paper by the same group showed that Rab11 is essential in controlling the activation of RTKs at the leading edge of these cells [76]. Both overexpression and depletion of *Drosophila* Evi5 abolished enrichment of phosphotyrosine signals at the front of the cell cluster, indicating that stringent regulation of Rab11 activity is necessary for the proper spatial restriction of activated signaling receptors [75]. Acting as a Rab GAP of the Rab11 family therefore, Evi5 plays clear and important roles in modulating cytokinesis and cell migration during development.

5. Evi5 in human diseases

The association of Evi5 with important cell cycle and cell migration regulators suggests its involvement in the pathology of certain diseases. Early work associated with Evi5's identification points towards its role as an oncogene or tumor suppressor [2,3]. This notion is strengthened by its interactions with Emi1 and Rab11, with an apparent role in the regulation of the cell cycle through interaction with the aurora-B kinase protein complex [4,12]. However, direct evidence for Evi5's role in oncogenesis, or suppression

of tumorigenesis, is still lacking. Evi5 has been found to be overexpressed in certain cancer types [77,78]. For instance, Evi5 could serve to prevent exhaustion of hematopoietic stem cells, whose levels are elevated as a result of the loss of tumor suppressor Runt-related transcription factor 1 (RUNX1) [78]. This suggests that elevated Evi5 levels could promote leukemia in cooperation with a loss of RUNX1. It is also conceivable that loss of Evi5 may enhance cancer cell invasion and metastasis via an enhanced Rab11 family GTPase activity, but this possibility remains to be clearly demonstrated.

An unexpected Evi5 disease link has arose from recent genomic analyses, which implicated Evi5 as a risk factor for multiple sclerosis (MS), a rather common autoimmune demyelinating disease with multiple neurological symptoms [79]. MS has a modest degree of heritability, and the Human Leukocyte Antigen (HLA) region is a known strong susceptibility locus for MS [80]. Genome-wide association studies have since identified other susceptibility genes, including *CD25*, *CD58*, *CLEC16a*, Interleukin-2 Receptor α (*IL2RA*), Interleukin-7 Receptor (*IL7R*), *G5* (*GPC5*), Regulator of G protein Signaling 1 (*RGS1*), and Tyrosine Kinase 2 (*TYK2*) [81–83], with some of these genes having clear functions in T-cell-mediated immunity.

Several separate analyses have associated single nucleotide polymorphisms (SNPs) of Evi5 with increased MS susceptibility [82,84–89]. The molecular basis of this association is unclear, as a role for Evi5 in T-cell mediated immunity has not yet been explored extensively. However, it is conceivable that Evi5 could affect aspects of innate immunity such as toll-like receptor signaling through Rab11, which was shown to be responsible for transport of the receptor via recycling endosomes [57]. A recent analysis of enhancer elements bound by the CCCTC-binding protein (CTCF) indicated that MS susceptibility SNPs within the last intron of




Fig. 3. Evi5's role in cytokinesis. Rab11a (Rab11) and Evi5 have both been localized to the pericentriolar region in U2OS cells (the cross indicated as 'C' in the diagram). Rab11a, together with its effectors (Rab11-FIP), are required to target recycling endosomes to the midbody or cleavage furrow, carrying physical membrane and factors important for cytokinesis. Evi5's presence at the interphase centrosomes and recycling endosomes could inhibit FIP's promotion of cytokinesis [42]. It was also noted that in prophase, Evi5 appears to have a cytoplasmic localization. However in metaphase, Evi5 become associated with the mitotic spindles and it moves to the spindle midzone and midbody at the final stages of mitosis. At these late stages Evi5 also interacts with the Chromosomal Passenger Complex (CPC) INCENP-Aurora kinase B (AURKB)-survivin (S) for the completion of cytokinesis [12]. From the different views on Evi5's localization, it is possible that Evi5 plays not one, but possibly several roles in cell division, with the differing roles depend critically on its spatial or temporal localization (that are in turn regulated by ubiquitination and degradation).

the *Evi5* gene, likely CTCF-binding enhancer elements, affected the expression of the neighboring gene and not *Evi5* itself [89]. That *Gfi1* and not *Evi5* being the causal gene for MS susceptibility, at least pertaining to some of the SNPs, makes intuitive sense, as *Gfi1* is a known regulator of lymphocyte development and activation [90].

6. Functional roles of *Evi5*-like (*Evi5L*)

While *Evi5* has been implicated in a variety of roles and several diseases, the function of *Evi5L*, a paralogue of *Evi5*, is more elusive. It bears approximately 70% homology to *Evi5*, and hence would be expected to interact to some degree with *Evi5* interacting partners, yet, no interaction has been demonstrated. Unlike *Evi5*, *Evi5L* appears to be largely cytosolic throughout the cell cycle, but has an interesting apparent concentration at the pericentriolar region (Fig. 4). *Evi5L* was first shown to bind Rab10 and exhibits in vitro GAP activity towards the small GTPase [5]. Rab10 is important in the regulation of endocytic recycling, and has been implicated in multiple roles that include basolateral recycling of clathrin-independent cargo in *Caenorhabditis elegans* intestine [91], AMPA-type glutamate receptors in post-synaptic membranes [92], basolateral recycling in polarized Madin–Darby canine kidney (MDCK) cells, Glut4 transporter trafficking in response to insulin [93], Toll-like receptor 4 (TLR4) signaling [94], phagosome maturation [95], and ciliary transport [96]. Notably, except for primary ciliogenesis, the role of *Evi5L* in these processes has not yet been demonstrated.

The primary cilium is a singular cellular projection protruding from the surface of cells that concentrates various signaling receptors and molecules. Hence, it is considered to be a signaling nexus that plays key roles in cellular physiology and organism development [97,98]. In fact, defects in the primary cilium's functional and trafficking components underlie multiple human ciliopathies [99]. Screening for Rab GAPs that affect primary cilium formation, Barr and colleagues identified Rab8a, –17, and –23, and their cognate GAPs, XM_037557, TBC1D7, and *Evi5L*. Carried out in the human telomerase-immortalized retinal pigmented epithelial (hTERT-RPE1) cell model, cells overexpressing *Evi5L* significantly

reduced primary cilium formation by more than half [100]. In the in vitro GAP activity assays, the authors showed that *Evi5L* especially exhibited specific activity towards Rab23, but not Rab10.

Rab23 was first isolated based on a mouse mutation to a gene whose functions antagonized Sonic hedgehog (Shh) signaling during embryonic development [101]. Shh signaling is critically dependent on the primary cilia, as proper functioning of ciliary intraflagellar transport (IFT) processes are essential for Shh signaling [102,103], and key components of the Shh signaling pathway, such as the Gli family of transcription factors and smoothened, are enriched at the primary cilia [104,105]. Although Rab23 has not yet been shown to be critical for ciliogenesis and ciliary transport as shown for Rab8a [106,107] and Rab11 [66], it is known to influence the levels of smoothened in the cilium [108]. Notably, in humans, Rab23 mutations underlie the pathological phenotypes associated with Carpenter's syndrome [109], whose phenotypes bear resemblance to those found in some ciliopathies, especially obesity, polydactyly and certain malformations of the brain [109]. Although the exact ciliary cargo involved and transport step mediated by Rab23 and *Evi5L* are yet unclear, accumulating evidence suggest that they have important roles in specific ciliary transport.

7. Future perspectives

We have summarized and discussed above cellular and physiological activities and roles of the *Evi5* family of TBC proteins. The tumorigenic (or tumor suppressive) activity of *Evi5* requires further clarification. Its ability to prevent hematopoietic stem cell exhaustion [78] suggests that *Evi5* might have a more general role in adult stem cells. It is also not known if *Evi5* has a direct role in cancer cell migration and invasion via its interaction with Rab11GTPases and the Rab coupling protein (RCP) [110]. Many potential functional aspects of *Evi5* and *Evi5L* remain unexplored. One possibility pertains to a plausible role for *Evi5* in ciliary transport and ciliogenesis via its GAP activity on Rab11 [66]. A recent report showed that *Evi5* levels are dramatically elevated during blastema formation in salamander limb regeneration [111]. The




Fig. 4. Subcellular localization of *Evi5L*. Top panel: A magnified image of *Evi5L* in NIH3T3 mouse fibroblasts using an *Evi5L* antibody from Abcam (ab88915). Endogenous *Evi5L* (red) appears to have a generally diffused cytosolic staining in NIH3T3 mouse fibroblasts. Cells are transfected with Arl13b-GFP (Origene; MG206808) (green), a primary cilium marker, and also labeled with Pericentrin (Abcam; ab4448) (magenta). Bottom panel: A magnified image of an NIH3T3 mouse fibroblast that is transiently overexpressing both mouse Arl13b-GFP (green) and human *Evi5L* (Openbiosystems; MHS1011-76129), and co-stained with Pericentrin (Abcam; ab4448) (magenta) and *Evi5L* (ab88915) (red). In both panels of images, primary cilia were induced after treatment with 24 h of serum starvation media. From the panel of images, it appears that *Evi5L* staining exhibits a partial co-localization with the pericentrin staining. Scale bar = 5 μ m.

authors postulated that high levels of Evi5 function to arrest dedifferentiated cells in the G1/S/G2 phases of the cell cycle until they have accumulated under the wound epidermis, with entry into mitosis subsequently triggered by growth factors. It would be interesting to see if Evi5 levels are also similarly elevated during regenerative attempts in mammalian tissues, which we know have a vastly inferior regenerative capacity compared to amphibians. Similarly, it would also be interesting to investigate Evi5's role in cell migration during wound healing.

While activity of Evi5 may be restricted by its nuclear and centrosomal localization [11], Evi5L is largely cytosolic, with some concentration around the pericentriolar region (Fig. 3). It could conceivably participate in trafficking events in multiple membrane compartments. While its overexpression clearly affects primary ciliogenesis [100], its exact mode of function is unclear. Although Evi5L exhibits specific Rab23 GAP activity in vitro, Evi5L's role in Shh signaling has not been reported. Moreover, whether it functions strictly through Rab23 is unclear. Alternatively, its suspected in vitro interaction with Rab10, reported by one group [5] but not another, is worth further exploration. Thus far, Evi5 and Evi5L appears to have non-overlapping Rab targets and functions, it would be interesting to see if these could, at least in some cases, complement the deficiency of each other when overexpressed.

Investigating the possible underlying cellular mechanisms of how Evi5 polymorphisms influence onset and progression of diseases such as MS would be of translational importance. Current animal models of MS mimic disease pathology, but not its genesis. It would be worth checking if transgenic expression of Evi5 and Gfi1, or their specific ectopic expression in the central nervous system in animal models will either produce MS pathologies, or enhance the susceptibility to MS induced by neurochemical means.

Acknowledgements

The authors declare no conflict of interest. B.L.T. is supported by the NUS Graduate School of Integrative Sciences and Engineering. We are grateful to the constructive comments from both reviewers, which improved the manuscript.

References

- [1] Liao, X., Buchberg, A.M., Jenkins, N.A. and Copeland, N.G. (1995) Evi-5, a common site of retroviral integration in AKXD T-cell lymphomas, maps near Gfi-1 on mouse chromosome 5. *J. Virol.* 69, 7132–7137.
- [2] Liao, X., Du, Y., Morse, H.C., Jenkins, N.A. and Copeland, N.G. (1997) Proviral integrations at the Evi5 locus disrupt a novel 90 kDa protein with homology to the Tre2 oncogene and cell-cycle regulatory proteins. *Oncogene* 14, 1023–1029.
- [3] Roberts, T., Chernova, O. and Cowell, J.K. (1998) NB4S, a member of the TBC1 domain family of genes, is truncated as a result of a constitutional t(1;10)(p22;q21) chromosome translocation in a patient with stage 4S neuroblastoma. *Hum. Mol. Genet.* 7, 1169–1178.
- [4] Eldridge, A.G., Loktev, A.V., Hansen, D.V., Verschuren, E.W., Reimann, J.D.R. and Jackson, P.K. (2006) The Evi5 oncogene regulates cyclin accumulation by stabilizing the anaphase-promoting complex inhibitor Emi1. *Cell* 124, 367–380.
- [5] Itoh, T., Satoh, M., Kanno, E. and Fukuda, M. (2006) Screening for target Rabs of TBC (Tre-2/Bub2/Cdc16) domain-containing proteins based on their Rab-binding activity. *Genes Cells* 11, 1023–1037.
- [6] Frasa, M.A.M., Koessmeier, K.T., Ahmadian, M.R. and Braga, V.M.M. (2012) Illuminating the functional and structural repertoire of human TBC/RABGAPs. *Nat. Rev. Mol. Cell Biol.* 13, 67–73.
- [7] Hirano, T. (2006) At the heart of the chromosome: SMC proteins in action. *Nat. Rev. Mol. Cell Biol.* 7, 311–322.
- [8] Fukuda, M. (2011) TBC proteins: GAPs for mammalian small GTPase Rab? *Biosci. Rep.* 31, 159–168.
- [9] Graumann, P.L. and Knust, T. (2009) Dynamics of the bacterial SMC complex and SMC-like proteins involved in DNA repair. *Chromosome Res.* 17, 265–275.
- [10] Wood, A.J., Severson, A.F. and Meyer, B.J. (2010) Condensin and cohesin complexity: the expanding repertoire of functions. *Nat. Rev. Genet.* 11, 391–404.
- [11] Fatar, S.L., Dabbeek, J.T.S., Ranalli, T.A. and Cowell, J.K. (2005) EVI5 is a novel centrosomal protein that binds to alpha- and gamma-tubulin. *Genomics* 86, 594–605.
- [12] Fatar, S.L., Sossey-Alaoui, K., Ranalli, T.A. and Cowell, J.K. (2006) EVI5 protein associates with the INCENP-aurora B kinase-survivin chromosomal passenger complex and is involved in the completion of cytokinesis. *Exp. Cell Res.* 312, 2325–2335.
- [13] van der Waal, M.S., Hengeveld, R.C.C., van der Horst, A. and Lens, S.M.A. (2012) Cell division control by the chromosomal passenger complex. *Exp. Cell Res.* 318, 1407–1420.
- [14] Capalbo, L., Montebault, E., Takeda, T., Bassi, Z.L., Glover, D.M. and D'Avino, P.P. (2012) The chromosomal passenger complex controls the function of endosomal sorting complex required for transport-III Snf7 proteins during cytokinesis. *Open Biol.* 2, 120070.
- [15] Fielding, A.B., Schonteich, E., Matheson, J., Wilson, G., Yu, X., Hickson, G.R.X., Srivastava, S., Baldwin, S.A., Prekeris, R. and Gould, G.W. (2005) Rab11-FIP3 and FIP4 interact with Arf6 and the exocyst to control membrane traffic in cytokinesis. *EMBO J.* 24, 3389–3399.
- [16] Wilson, G.M., Fielding, A.B., Simon, G.C., Yu, X., Andrews, P.D., Hames, R.S., Frey, A.M., Peden, A.A., Gould, G.W. and Prekeris, R. (2005) The FIP3–Rab11 protein complex regulates recycling endosome targeting to the cleavage furrow during late cytokinesis. *Mol. Biol. Cell* 16, 849–860.
- [17] Reimann, J.D., Freed, E., Hsu, J.Y., Kramer, E.R., Peters, J.M. and Jackson, P.K. (2001) Emi1 is a mitotic regulator that interacts with Cdc20 and inhibits the anaphase promoting complex. *Cell* 105, 645–655.
- [18] Hsu, J.Y., Reimann, J.D.R., Sørensen, C.S., Lukas, J. and Jackson, P.K. (2002) E2F-dependent accumulation of hEmi1 regulates S phase entry by inhibiting APC(Cdh1). *Nat. Cell Biol.* 4, 358–366.
- [19] Hansen, D.V., Loktev, A.V., Ban, K.H. and Jackson, P.K. (2004) Plk1 regulates activation of the anaphase promoting complex by phosphorylating and triggering SCFbetaTrCP-dependent destruction of the APC Inhibitor Emi1. *Mol. Biol. Cell* 15, 5623–5634.
- [20] Machida, Y.J. and Dutta, A. (2007) The APC/C inhibitor, Emi1, is essential for prevention of rereplication. *Genes Dev.* 21, 184–194.
- [21] Margottin-Goguet, F., Hsu, J.Y., Loktev, A., Hsieh, H.M., Reimann, J.D.R. and Jackson, P.K. (2003) Prophase destruction of Emi1 by the SCF(betaTrCP/Slimb) ubiquitin ligase activates the anaphase promoting complex to allow progression beyond prometaphase. *Dev. Cell* 4, 813–826.
- [22] Stenmark, H. (2009) Rab GTPases as coordinators of vesicle traffic. *Nat. Rev. Mol. Cell Biol.* 10, 513–525.
- [23] Hutagalung, A.H. and Novick, P.J. (2011) Role of Rab GTPases in membrane traffic and cell physiology. *Physiol. Rev.* 91, 119–149.
- [24] Cai, H., Reinisch, K. and Ferro-Novick, S. (2007) Coats, tethers, Rabs, and SNAREs work together to mediate the intracellular destination of a transport vesicle. *Dev. Cell* 12, 671–682.
- [25] Epp, N., Rethmeier, R., Krämer, L. and Ungermann, C. (2011) Membrane dynamics and fusion at late endosomes and vacuoles – Rab regulation, multisubunit tethering complexes and SNAREs. *Eur. J. Cell Biol.* 90, 779–785.
- [26] Angers, C.G. and Merz, A.J. (2011) New links between vesicle coats and Rab-mediated vesicle targeting. *Semin. Cell Dev. Biol.* 22, 18–26.
- [27] Horgan, C.P. and McCaffrey, M.W. (2011) Rab GTPases and microtubule motors. *Biochem. Soc. Trans.* 39, 1202–1206.
- [28] Barr, F. and Lambright, D.G. (2010) Rab GEFs and GAPs. *Curr. Opin. Cell Biol.* 22, 461–470.
- [29] Buday, L. and Downward, J. (2008) Many faces of Ras activation. *Biochim. Biophys. Acta* 1786, 178–187.
- [30] Delprat, A., Merithew, E. and Lambright, D.G. (2004) Structure, exchange determinants, and family-wide rab specificity of the tandem helical bundle and Vps9 domains of Rabex-5. *Cell* 118, 607–617.
- [31] Marat, A.L., Dokainish, H. and McPherson, P.S. (2011) DENN domain proteins: regulators of Rab GTPases. *J. Biol. Chem.* 286, 13791–13800.
- [32] Bernards, A. (2003) GAPs galore! A survey of putative Ras superfamily GTPase activating proteins in man and Drosophila. *Biochim. Biophys. Acta* 1603, 47–82.
- [33] Rivera-Molina, F.E. and Novick, P.J. (2009) A Rab GAP cascade defines the boundary between two Rab GTPases on the secretory pathway. *Proc. Natl. Acad. Sci. USA* 106, 14408–14413.
- [34] Richardson, P.M. and Zon, L.I. (1995) Molecular cloning of a cDNA with a novel domain present in the tre-2 oncogene and the yeast cell cycle regulators BUB2 and cdc16. *Oncogene* 11, 1139–1148.
- [35] Strom, M., Vollmer, P., Tan, T.J. and Gallwitz, D. (1993) A yeast GTPase-activating protein that interacts specifically with a member of the Ypt/Rab family. *Nature* 361, 736–739.
- [36] Cuif, M.H., Possmayer, F., Zander, H., Bordes, N., Jollivet, F., Couedel-Courteille, A., Janoueix-Lerosey, I., Langsley, G., Bornens, M. and Goud, B. (1999) Characterization of GAPCenA, a GTPase activating protein for Rab6, part of which associates with the centrosome. *EMBO J.* 18, 1772–1782.
- [37] Lanzetti, L., Rybin, V., Malabarba, M.G., Christoforidis, S., Scita, G., Zerial, M. and Di Fiore, P.P. (2000) The Eps8 protein coordinates EGF receptor signalling through Rac and trafficking through Rab5. *Nature* 408, 374–377.
- [38] Pei, L., Peng, Y., Yang, Y., Ling, X.B., Van Eynhoven, W.G., Nguyen, K.C.Q., Rubin, M., Hoey, T., Powers, S. and Li, J. (2002) PRC17, a novel oncogene encoding a Rab GTPase-activating protein, is amplified in prostate cancer. *Cancer Res.* 62, 5420–5424.

- [39] Haas, A.K., Fuchs, E., Kopajtich, R. and Barr, F.A. (2005) A GTPase-activating protein controls Rab5 function in endocytic trafficking. *Nat. Cell Biol.* 7, 887–893.
- [40] Miinea, C.P., Sano, H., Kane, S., Sano, E., Fukuda, M., Peränen, J., Lane, W.S. and Lienhard, G.E. (2005) AS160, the Akt substrate regulating GLUT4 translocation, has a functional Rab GTPase-activating protein domain. *Biochem. J.* 391, 87–93.
- [41] Kanno, E., Ishibashi, K., Kobayashi, H., Matsui, T., Ohbayashi, N. and Fukuda, M. (2010) Comprehensive screening for novel rab-binding proteins by GST pull-down assay using 60 different mammalian Rabs. *Traffic* 11, 491–507.
- [42] Westlake, C.J., Junutula, J.R., Simon, G.C., Pilli, M., Prekeris, R., Scheller, R.H., Jackson, P.K. and Eldridge, A.G. (2007) Identification of Rab11 as a small GTPase binding protein for the Evi5 oncogene. *Proc. Natl. Acad. Sci. USA* 104, 1236–1241.
- [43] Dabbeek, J.T.S., Faitar, S.L., Dufresne, C.P. and Cowell, J.K. (2007) The EVI5 TBC domain provides the GTPase-activating protein motif for RAB11. *Oncogene* 26, 2804–2808.
- [44] Horgan, C.P. and McCaffrey, M.W. (2009) The dynamic Rab11-FIPs. *Biochem. Soc. Trans.* 37, 1032–1036.
- [45] Ullrich, O., Reinsch, S., Urbé, S., Zerial, M. and Parton, R.G. (1996) Rab11 regulates recycling through the pericentriolar recycling endosome. *J. Cell Biol.* 135, 913–924.
- [46] Prekeris, R., Klumperman, J. and Scheller, R.H. (2000) A Rab11/Rip11 protein complex regulates apical membrane trafficking via recycling endosomes. *Mol. Cell* 6, 1437–1448.
- [47] Lindsay, A.J. and McCaffrey, M.W. (2002) Rab11-FIP2 functions in transferrin recycling and associates with endosomal membranes via its COOH-terminal domain. *J. Biol. Chem.* 277, 27193–27199.
- [48] Su, T., Bryant, D.M., Lutton, F., Vergés, M., Ulrich, S.M., Hansen, K.C., Datta, A., Eastburn, D.J., Burlingame, A.L., Shokat, K.M. and Mostov, K.E. (2010) A kinase cascade leading to Rab11-FIP5 controls transcytosis of the polymeric immunoglobulin receptor. *Nat. Cell Biol.* 12, 1143–1153.
- [49] McEwen, D.P., Schumacher, S.M., Li, Q., Benson, M.D., Iñiguez-Lluhi, J.A., Van Genderen, K.M. and Martens, J.R. (2007) Rab-GTPase-dependent endocytic recycling of Kv1.5 in atrial myocytes. *J. Biol. Chem.* 282, 29612–29620.
- [50] Silvius, M.R., Bertrand, C.A., Ameen, N., Golín-Bisello, F., Butterworth, M.B., Frizzell, R.A. and Bradbury, N.A. (2009) Rab11b regulates the apical recycling of the cystic fibrosis transmembrane conductance regulator in polarized intestinal epithelial cells. *Mol. Biol. Cell* 20, 2337–2350.
- [51] Butterworth, M.B., Edinger, R.S., Silvius, M.R., Gallo, L.L., Liang, X., Apodaca, G., Fizzell, R.A. and Johnson, J.P. (2012) Rab11b regulates the trafficking and recycling of the epithelial sodium channel (ENaC). *Am. J. Physiol. Renal Physiol.* 302, F581–F590.
- [52] Cullis, D.N., Philip, B., Baleja, J.D. and Feig, L.A. (2002) Rab11-FIP2, an adaptor protein connecting cellular components involved in internalization and recycling of epidermal growth factor receptors. *J. Biol. Chem.* 277, 49158–49166.
- [53] Moore, R.H., Millman, E.E., Alpizar-Foster, E., Dai, W. and Knoll, B.J. (2004) Rab11 regulates the recycling and lysosome targeting of beta2-adrenergic receptors. *J. Cell Sci.* 117, 3107–3117.
- [54] Parent, A., Hamelin, E., Germain, P. and Parent, J.L. (2009) Rab11 regulates the recycling of the beta2-adrenergic receptor through a direct interaction. *Biochem. J.* 418, 163–172.
- [55] Gardner, L.A., Hajjhussein, H., Frederick-Dyer, K.C. and Bahouth, S.W. (2011) Rab11a and its binding partners regulate the recycling of the β 1-adrenergic receptor. *Cell. Signal.* 23, 46–57.
- [56] Kumar, A., Kremer, K.N., Dominguez, D., Tadi, M. and Hedin, K.E. (2011) G α 13 and Rho mediate endosomal trafficking of CXCR4 into Rab11+ vesicles upon stromal cell-derived factor-1 stimulation. *J. Immunol.* 186, 951–958.
- [57] Kagan, J.C. (2010) Recycling endosomes and TLR signaling – the Rab11 GTPase leads the way. *Immunity* 33, 578–580.
- [58] Chi, S., Cao, H., Wang, Y. and McNiven, M.A. (2011) Recycling of the epidermal growth factor receptor is mediated by a novel form of the clathrin adaptor protein Eps15. *J. Biol. Chem.* 286, 35196–35208.
- [59] Lock, J.G. and Stow, J.L. (2005) Rab11 in recycling endosomes regulates the sorting and basolateral transport of E-cadherin. *Mol. Biol. Cell* 16, 1744–1755.
- [60] Caswell, P.T., Chan, M., Lindsay, A.J., McCaffrey, M.W., Boettiger, D. and Norman, J.C. (2008) Rab-coupling protein coordinates recycling of alpha5beta1 integrin and EGFR1 to promote cell migration in 3D microenvironments. *J. Cell Biol.* 183, 143–155.
- [61] Liu, J., Zhang, J.P., Shi, M., Quinn, T., Bradner, J., Beyer, R., Chen, S. and Zhang, J. (2009) Rab11a and HSP90 regulate recycling of extracellular alpha-synuclein. *J. Neurosci.* 29, 1480–1485.
- [62] Jing, J. and Prekeris, R. (2009) Polarized endocytic transport: the roles of Rab11 and Rab11-FIPs in regulating cell polarity. *Histol. Histopathol.* 24, 1171–1180.
- [63] Emery, G., Hutterer, A., Berdnik, D., Mayer, B., Wirtz-Peitz, F., Gaitan, M.G. and Knoblich, J.A. (2005) Asymmetric Rab 11 endosomes regulate delta recycling and specify cell fate in the Drosophila nervous system. *Cell* 122, 763–773.
- [64] Desclozeaux, M., Venturato, J., Wylie, F.G., Kay, J.G., Joseph, S.R., Le, H.T. and Stow, J.L. (2008) Active Rab11 and functional recycling endosome are required for E-cadherin trafficking and lumen formation during epithelial morphogenesis. *Am. J. Physiol. Cell Physiol.* 295, C545–C556.
- [65] Mateus, A.M., Gorfinkiel, N., Schamberg, S. and Martinez Arias, A. (2011) Endocytic and recycling endosomes modulate cell shape changes and tissue behavior during morphogenesis in Drosophila. *PLoS ONE* 6, e18729.
- [66] Knödler, A., Feng, S., Zhang, J., Zhang, X., Das, A., Peränen, J. and Guo, W. (2010) Coordination of Rab8 and Rab11 in primary ciliogenesis. *Proc. Natl. Acad. Sci. USA* 107, 6346–6351.
- [67] Longatti, A., Lamb, C.A., Razi, M., Yoshimura, S.I., Barr, F.A. and Tooze, S.A. (2012) TBC1D14 regulates autophagosome formation via Rab11- and ULK1-positive recycling endosomes. *J. Cell Biol.* 197, 659–675.
- [68] Giansanti, M.G., Belloni, G. and Gatti, M. (2007) Rab11 is required for membrane trafficking and actomyosin ring constriction in meiotic cytokinesis of Drosophila males. *Mol. Biol. Cell* 18, 5034–5047.
- [69] Simon, G.C. and Prekeris, R. (2008) The role of FIP3-dependent endosome transport during cytokinesis. *Commun. Integr. Biol.* 1, 132–133.
- [70] Kawauchi, T., Sekine, K., Shikanai, M., Chihama, K., Tomita, K., Kubo, K.I., Nakajima, K., Nabeshima, Y.I. and Hoshino, M. (2010) Rab GTPases-dependent endocytic pathways regulate neuronal migration and maturation through N-cadherin trafficking. *Neuron* 67, 588–602.
- [71] Kessler, D., Gruen, G.C., Heider, D., Morgner, J., Reis, H., Schmid, K.W. and Jendrosseck, V. (2012) The action of small GTPases Rab11 and Rab25 in vesicle trafficking during cell migration. *Cell. Physiol. Biochem.* 29, 647–656.
- [72] Li, X., Sapp, E., Chase, K., Comer-Tierney, L.A., Masso, N., Alexander, J., Reeves, P., Kegel, K.B., Valencia, A., Esteves, M., Aronin, N. and Difiglia, M. (2009) Disruption of Rab11 activity in a knock-in mouse model of Huntington's disease. *Neurobiol. Dis.* 36, 374–383.
- [73] Richards, P., Didszun, C., Campesan, S., Simpson, A., Horley, B., Young, K.W., Glynn, P., Cain, K., Kyriacou, C.P., Giorgini, F. and Nicotera, P. (2011) Dendritic spine loss and neurodegeneration is rescued by Rab11 in models of Huntington's disease. *Cell Death Differ.* 18, 191–200.
- [74] Caswell, P.T., Spence, H.J., Parsons, M., White, D.P., Clark, K., Cheng, K.W., Mills, G.B., Humphries, M.J., Messent, A.J., Anderson, K.I., McCaffrey, M.W., Ozanne, B.W. and Norman, J.C. (2007) Rab25 associates with alpha5beta1 integrin to promote invasive migration in 3D microenvironments. *Dev. Cell* 13, 496–510.
- [75] Laflamme, C., Assaker, G., Ramel, D., Dorn, J.F., She, D., Maddox, P.S. and Emery, G. (2012) Evi5 promotes collective cell migration through its Rab-GAP activity. *J. Cell Biol.* 198, 57–67.
- [76] Assaker, G., Ramel, D., Wculek, S.K., González-Gaitán, M. and Emery, G. (2010) Spatial restriction of receptor tyrosine kinase activity through a polarized endocytic cycle controls border cell migration. *Proc. Natl. Acad. Sci. USA* 107, 22558–22563.
- [77] Varghese, S., Xu, H., Bartlett, D., Hughes, M., Pingpank, J.F., Beresnev, T. and Alexander, H.R. (2010) Isolated hepatic perfusion with high-dose melphalan results in immediate alterations in tumor gene expression in patients with metastatic ocular melanoma. *Ann. Surg. Oncol.* 17, 1870–1877.
- [78] Jacob, B., Osato, M., Yamashita, N., Wang, C.Q., Taniuchi, I., Littman, D.R., Asou, N. and Ito, Y. (2010) Stem cell exhaustion due to Runx1 deficiency is prevented by Evi5 activation in leukemogenesis. *Blood* 115, 1610–1620.
- [79] Compston, A. and Coles, A. (2002) Multiple sclerosis. *Lancet* 359, 1221–1231.
- [80] Oksenberg, J.R., Baranzini, S.E., Sawcer, S. and Hauser, S.L. (2008) The genetics of multiple sclerosis: SNPs to pathways to pathogenesis. *Nat. Rev. Genet.* 9, 516–526.
- [81] Wellcome Trust Case Control Consortium et al. (2007) Association scan of 14,500 non-synonymous SNPs in four diseases identifies autoimmunity variants. *Nat. Genet.* 39, 1329–1337.
- [82] International Multiple Sclerosis Genetics Consortium, Hafler, D.A., Compston, A., Sawcer, S., Lander, E.S., Daly, M.J., De Jager, P.L., de Bakker, P.I.W., Gabriel, S.B., Mirel, D.B., Ivinson, A.J., Pericak-Vance, M.A., Gregory, S.G., Rioux, J.D., McCauley, J.L., Haines, J.L., Barcellos, L.F., Cree, B., Oksenberg, J.R. and Hauser, S.L. (2007) Risk alleles for multiple sclerosis identified by a genome-wide study. *N. Engl. J. Med.* 357, 851–862.
- [83] Baranzini, S.E. et al. (2009) Genome-wide association analysis of susceptibility and clinical phenotype in multiple sclerosis. *Hum. Mol. Genet.* 18, 767–778.
- [84] Hoppenbrouwers, I.A., Aulchenko, Y.S., Ebers, G.C., Ramagopalan, S.V., Oostra, B.A., van Duijn, C.M. and Hintzen, R.Q. (2008) EVI5 is a risk gene for multiple sclerosis. *Genes Immun.* 9, 334–337.
- [85] D'Netto, M.J., Ward, H., Morrison, K.M., Ramagopalan, S.V., Dymmet, D.A., DeLuca, G.C., Handunnetthi, L., Sadovnick, A.D. and Ebers, G.C. (2009) Risk alleles for multiple sclerosis in multiplex families. *Neurology* 72, 1984–1988.
- [86] Australia and New Zealand Multiple Sclerosis Genetics Consortium (ANZgene) (2009) Genome-wide association study identifies new multiple sclerosis susceptibility loci on chromosomes 12 and 20. *Nat. Genet.* 41, 824–828.
- [87] Alcina, A., Fernández, O., Gonzalez, J.R., Catalá-Rabasa, A., Fedetz, M., Ndagire, D., Leyva, L., Guerrero, M., Arnal, C., Delgado, C., Lucas, M., Izquierdo, G. and Matesanz, F. (2010) Tag-SNP analysis of the GF11-EVI5-RPL5-FAM69 risk locus for multiple sclerosis. *Eur. J. Hum. Genet.* 18, 827–831.
- [88] Johnson, B.A., Wang, J., Taylor, E.M., Caillier, S.J., Herbert, J., Khan, O.A., Cross, A.H., De Jager, P.L., Gourraud, P.A.F., Cree, B.C.A., Hauser, S.L. and Oksenberg, J.R. (2010) Multiple sclerosis susceptibility alleles in African Americans. *Genes Immun.* 11, 343–350.
- [89] Martin, D. et al. (2011) Genome-wide CTCF distribution in vertebrates defines equivalent sites that aid the identification of disease-associated genes. *Nat. Struct. Mol. Biol.* 18, 708–714.

- [90] Möröy, T. and Khandanpour, C. (2011) Growth factor independence 1 (Gfi1) as a regulator of lymphocyte development and activation. *Semin. Immunol.* 23, 368–378.
- [91] Shi, A., Chen, C.C.H., Banerjee, R., Glodowski, D., Audhya, A., Rongo, C. and Grant, B.D. (2010) EHBP-1 functions with RAB-10 during endocytic recycling in *Caenorhabditis elegans*. *Mol. Biol. Cell* 21, 2930–2943.
- [92] Glodowski, D.R., Chen, C.C.H., Schaefer, H., Grant, B.D. and Rongo, C. (2007) RAB-10 regulates glutamate receptor recycling in a cholesterol-dependent endocytosis pathway. *Mol. Biol. Cell* 18, 4387–4396.
- [93] Sano, H., Roach, W.G., Peck, G.R., Fukuda, M. and Lienhard, G.E. (2008) Rab10 in insulin-stimulated GLUT4 translocation. *Biochem. J.* 411, 89–95.
- [94] Wang, D., Lou, J., Ouyang, C., Chen, W., Liu, Y., Liu, X., Cao, X., Wang, J. and Lu, L. (2010) Ras-related protein Rab10 facilitates TLR4 signaling by promoting replenishment of TLR4 onto the plasma membrane. *Proc. Natl. Acad. Sci. USA* 107, 13806–13811.
- [95] Cardoso, C.M.P., Jordao, L. and Vieira, O.V. (2010) Rab10 regulates phagosome maturation and its overexpression rescues Mycobacterium-containing phagosomes maturation. *Traffic* 11, 221–235.
- [96] Babbey, C.M., Bacallao, R.L. and Dunn, K.W. (2010) Rab10 associates with primary cilia and the exocyst complex in renal epithelial cells. *Am. J. Physiol. Renal Physiol.* 299, F495–F506.
- [97] Goetz, S.C. and Anderson, K.V. (2010) The primary cilium: a signalling centre during vertebrate development. *Nat. Rev. Genet.* 11, 331–344.
- [98] Oh, E.C. and Katsanis, N. (2012) Cilia in vertebrate development and disease. *Development* 139, 443–448.
- [99] Sharma, N., Berbari, N.F. and Yoder, B.K. (2008) Ciliary dysfunction in developmental abnormalities and diseases. *Curr. Top. Dev. Biol.* 85, 371–427.
- [100] Yoshimura, S.I., Egerer, J., Fuchs, E., Haas, A.K. and Barr, F.A. (2007) Functional dissection of Rab GTPases involved in primary cilium formation. *J. Cell Biol.* 178, 363–369.
- [101] Eggenschwiler, J.T., Espinoza, E. and Anderson, K.V. (2001) Rab23 is an essential negative regulator of the mouse Sonic hedgehog signalling pathway. *Nature* 412, 194–198.
- [102] Liu, A., Wang, B. and Niswander, L.A. (2005) Mouse intraflagellar transport proteins regulate both the activator and repressor functions of Gli transcription factors. *Development* 132, 3103–3111.
- [103] Wong, S.Y. and Reiter, J.F. (2008) The primary cilium at the crossroads of mammalian hedgehog signaling. *Curr. Top. Dev. Biol.* 85, 225–260.
- [104] Haycraft, C.J., Banizs, B., Aydin-Son, Y., Zhang, Q., Michaud, E.J. and Yoder, B.K. (2005) Gli2 and Gli3 localize to cilia and require the intraflagellar transport protein polaris for processing and function. *PLoS Genet.* 1, e53.
- [105] Corbit, K.C., Aanstad, P., Singla, V., Norman, A.R., Stainier, D.Y.R. and Reiter, J.F. (2005) Vertebrate smoothened functions at the primary cilium. *Nature* 437, 1018–1021.
- [106] Nachury, M.V., Loktev, A.V., Zhang, Q., Westlake, C.J., Peränen, J., Merdes, A., Slusarski, D.C., Scheller, R.H., Bazan, J.F., Sheffield, V.C. and Jackson, P.K. (2007) A core complex of BBS proteins cooperates with the GTPase Rab8 to promote ciliary membrane biogenesis. *Cell* 129, 1201–1213.
- [107] Feng, S., Knödler, A., Ren, J., Zhang, J., Zhang, X., Hong, Y., Huang, S., Peränen, J. and Guo, W. (2012) A Rab8 guanine nucleotide exchange factor–effector interaction network regulates primary ciliogenesis. *J. Biol. Chem.* 287, 15602–15609.
- [108] Boehlke, C., Bashkurov, M., Buescher, A., Krick, T., John, A.K., Nitschke, R., Walz, G. and Kuehn, E.W. (2010) Differential role of Rab proteins in ciliary trafficking: Rab23 regulates smoothened levels. *J. Cell Sci.* 123, 1460–1467.
- [109] Jenkins, D., Seelow, D., Jehee, F.S., Perlyn, C.A., Alonso, L.G., Bueno, D.F., Donnai, D., Josifova, D., Josifiova, D., Mathijssen, I.M.J., Morton, J.E.V., Orstavik, K.H., Sweeney, E., Wall, S.A., Marsh, J.L., Nurnberg, P., Passos-Bueno, M.R. and Wilkie, A.O.M. (2007) RAB23 mutations in Carpenter syndrome imply an unexpected role for hedgehog signaling in cranial-suture development and obesity. *Am. J. Hum. Genet.* 80, 1162–1170.
- [110] Tang, B.L. (2010) Is Rab25 a tumor promoter or suppressor – context dependency on RCP status? *Tumour Biol.* 31, 359–361.
- [111] Rao, N., Jhamb, D., Milner, D.J., Li, B., Song, F., Wang, M., Voss, S.R., Palakal, M., King, M.W., Saranjami, B., Nye, H.L.D., Cameron, J.A. and Stocum, D.L. (2009) Proteomic analysis of blastema formation in regenerating axolotl limbs. *BMC Biol.* 7, 83.